Abstract

The purpose of the present study is to summarise the mechanisms of action of Spirulina, phytochemical, food safety. nanofiber scaffolds, PHB, highlight the potential effects of this alga on humans, and address current and potential future clinical applications, based primarily on in vivo studies and a few carefully designed in vitro studies, as well as the highest levels of evidence currently available in the literature. The objective of this study is to examine the antioxidant activity of new commercial products containing fresh Spirulina and the factors influencing it. Antioxidant activity varies depending on the combination of elements in the meal, according to several studies. government organisations growing Spirulina to reduce the cost of purchasing it from private businesses and supplying it as food to anganwadis and ashrams so that it can gradually eradicate the problem of malnutrition in India. The synergy coefficient was used to identify and express synergistic, antagonistic, and additive interactions between samples in order to gradually alleviate the issue of malnutrition in India and other nations. In the present composition review the bioactive Spirulina and potential for different foods applications in functional covered.

Keywords: Phytochemical, Food safety, Nanofiber scaffolds, PHB

Authors

Shah Rehana Anjum A

Assistant Professor Department of Botany Rizvi Education Society's Rizvi College of Arts, Science and Commerce Mumbai, Maharashtra, India shahrehanaa@yahoo.com

Firdaus M. Quraishi

Assistant Professor Department of Biotechnology Elphinstone College Mumbai, Mahatashtra, India

Hina Alim

Assistant Professor Department of Life Sciences University of Mumbai Mumbai, Maharashtra, India,

I. INTRODUCTION

Spirulina are multicellular and filamentous blue-green microalgae belonging to two separate genera Spirulina and Arthrospira and consists of about 15 species. Of these, Arthrospira platensis is the most common and widely available Spirulina and most of the published research and public health decision refers to this specific species. Spirulina is natural food belongs to plantae kingdom which consists of different phytochemicals. (Balakrishnan et al., 2009 and Dawes, 1998). This phytochemicals are biologically significant and plays a vital role in medicinal applications. Many studies revealed that phytochemicals from Spirulina used in cancer, tuberculosis, inflammation and many other blood related diseases (Shyamala et al., 2013). The bioactive compound includes proteins, carbohydrates, phenolic compounds, vitamins, and minerals. The production of bioactive compound is depends on environmental conditions such as temperature, pH, nutrients, metal ions and other chemicals (Wijesekara I. et al., 2011). These phytochemicals play an important role in the treatment of many chronic diseases. There are two main methods for the analysis of phytochemical screening such as qualitative and quantitative analysis. The qualitative tests are used to identify the constituents. (Wijesekara et al., 2011). Spirulina is marketed and consumed in several countries, including, U.S.A, Thailand, Taiwan, Vietnam, China, India and Cuba (Shanmugam M et al., 2000).

II. FIELD OF APPLICATION

Asia has used Spirulina for more than a millennium. It has been marketed as a food supplement due to the high-quality protein and additional beneficial components (Ali and Saleh, 2012, Borowitzka, 2013, Priyadarshani and Rath, 2012). Spirulina has recently gained popularity as a non-toxic, non-carcinogenic natural blue colourant for use in food and cosmetics (Pandey et al., 2013). In the US, the FDA has designated Spirulina extract as a colour additive exempt from certification and authorised its use in ready-to eat cereals (apart from extruded cereals), frostings, ice cream and frozen desserts, dessert coatings and toppings, beverage mixes and powders, yoghurts, custards, puddings, cottage cheese, gelatin, and breadcrumbs. Spirulina is a suitable dye for dairy goods, ice cream, and sweets including chewing gum, coatings, chocolate, sugar embellishments, and candies. The shelf life of the finished goods is increased during processing and storage by low temperatures and light exclusion. The colourant is unsuitable for commercial use for watery food items with pH values below 4.5 and alcohol contents above 20%. (Newsome et al., 2014). Additionally, Spirulina is utilised as an antioxidant, thickening, and water binding agent, especially in face, skin, and hair care products, as well as a pigment in cosmetics including lip Sticks, eyeliners, and eye shadows. (Priyadarshani and Rath, 2012). Numerous toxicological investigations have validated the safety of Spirulina. Spirulina is currently one of the items on the US Food and Drug Administration's list of item classified as Generally Recognised as Safe (GRAS). (Tarantino LM.2003.) (Belay A. 2003.) Spirulina biomass demonstrates antiallergenic, antibacterial, antifungal, anti-inflammatory antioxidant, and immunomodulating effects in addition to stimulating vital biological processes (Khan et al., 2005). As a result, the biomass of Spirulina LEB 18 added to scaffolds promotes cell proliferation and tissue regeneration (Morais et al., 2010. Qureshi MA et al., 1995).

III.EXPECTED OUTCOMES

With the efforts of researchers and the government, the potential of *Spirulina* must be noticed and brought into use for solving various problems like malnutrition on national and international levels.

Spirulina proves to be an excellent source of micro nutrients like iron iodine, vitamin a foliate zinc, vitamin B1 vitamin B2 and vitamin B3 as all these micronutrients happen to be very effective in recovery from malnutrition. Eradication of malnutrition in Maharashtra gradually with the help of *Spirulina* supplements manufactured by the government.

- Use of *Spirulina* ruminants or expired supplements as a natural fertilizer to crops and other agricultural produce.
- Use of cheaper spiral in a based supplements produced by government for people suffering from anaemia diabetes cholesterol and blood pressure related issues as it aids them very well.
- Eradication of malnutrition in Maharashtra gradually with the help of *Spirulina* supplements manufactured by the government.
- Approximately 3000 tons dry weight is currently produced annually in the United States, Thailand, India, Taiwan, China, Pakistan, and Burma (Raja et al., 2007)

IV. SCOPE APPROACH

The *Spirulina* Market is expected to reach \$1.1 billion by 2030, at a CAGR of 9.4% from 2023 to 2030, while in terms of volume, the market is expected to reach 102,381.3 tons by 2030, at a CAGR of 8% from 2023 to 2030

V. USE OF SPIRULINA FOOD INDUSTRY

Spirulina are multicellular and filamentous blue-green algae that has gained considerable popularity in the health food industry and increasingly as a protein and vitamin supplement to aquaculture diets.

Earthrise farm was the first *Spirulina* farm in North America, established in 1976, and today is the largest *Spirulina* farm in the world. With over 40 years of experience and a 108-acre site, Earthrise has continued to produce high quality and safe *Spirulina* for consumers around the world. (WWW.Earthrise)

VI. BENEFITS TO THE SOCIETY

Like other blue-green algae, *Spirulina* is susceptible to contamination from poisonous compounds known as microcystins. Additionally, it can take up heavy metals from the water in which it is grown. For these reasons, it's crucial to purchase *Spirulina* from a reputable company.

Spirulina may strengthen the immune system, assist in preventing allergic reactions, and have antiviral and anticancer characteristics, according to test tube and animal research. However, there is no evidence that *Spirulina* offers these or any other advantages to humans. More study is required.

IIP Series, Volume 3, Book 14, Part 1, Chapter 9

SPIRULINA AS A SOURCE OF BIOACTIVE COMPOUNDS IN THE FOOD AND HEALTH SECTOR

Nanofiber scaffolds offer tissue engineering potential by replicating extracellular matrix structure and function. Electrospun scaffolds have high surface area, porosity, nutrient diffusion, and promote angiogenesis/vascularization during regeneration. (Ramier J, 2014)

Nanofiber scaffolds can be replaced with *Spirulina* biopolymers, which are biodegradable and compatible with cells and tissues. *Spirulina* biomass can be added to polymer solutions without extreme temperatures or pH, allowing internal components like proteins, fatty acids, and biopolymers to stimulate cells or tissues.

Polyhydroxyalkanoates (PHAs), including polyhydroxybutyrate (PHB), can be extracted from microorganisms like *Spirulina* and used as a toxic biocompatible scaffold for human tissue and organ culture. PHB degrades into a harmless compound, indicating its biocompatibility with cultured cells and tissues. *Spirulina* biopolymer nanofiber scaffolds have lower rejection risks and advantageous bioactive compounds (Sudesh K,et al 2000"and Jau M *et al.*, 2005).

VII.FUTURE SCOPE

Commercial and mass cultivation: In the early 1960s, Japan began a large-scale culture production programme for the microalgae chlorella and *Spirulina*. More than 22 nations currently engage in extensive commercial *Spirulina* cultivation.

VIII. CONCLUSION

Already, *Spirulina* is a well acknowledged and well-liked nutritional supplement. The requirement for studies to support its health advantages is minimal. This is supported by information on its nutritional worth, which makes *Spirulina* a superior option when creating diets and addressing malnutrition.

Preclinical research employing animal models is still being done to assess the bioactive potential of *Spirulina*. However, these researches appear to support the existence of *Spirulina's* potent antioxidant, anticancer, and antiviral activities as well as its ability to prevent diabetes, obesity, and allergic inflammation. Additionally, it has excellent hypoglycaemic, hypocholesterolaemia, and immunomodulatory potential.

The potential of *Spirulina* as a therapeutic food is drawing more attention due to consistent results on its bioactivities. The use of nutraceuticals in the treatment of hypercholesterolemia, hyperglycaemia, cardiovascular disease, and cancer should be quite beneficial. Studying the phytochemical composition, nutritional and biological effects allows the development of new scientific research.

REFERENCES

- [1] Shanmugam M, Mody KH. (2000) Heparinoid-active sulfated polysaccharides from marine algae as potential blood anticoagulant agents. Current Science; 79:1672-1683
- [2] Balakrishnan CP, Venkataraman K, Mohan VR, Louis JL, Athiperumal ST (2009). A general survey of the common agarophytes in the Gulf of Mannar in relation to agar ecology. Seaweed Research and Utilisation 2009; 31(1&2):33–46.
- [3] Dawes, C. (1998). Marine Botany. New York: John Wiley and Sons, Inc, 480.

- [4] Shyamala, V and N. Thangaraju. (2013) "Screening of Phytochemical and Antibacteriial activity of three different seaweeds from Gulf of Mannar, Tamil nadu". Phykos., Vol. 43(1), pp. 32-38.
- [5] Wijesekara I, Pangestuti R, Kim SK. (2011). Biological activities and potential health benefits of sulfated polysaccharides derived from marine algae. Carb Polymer. 84: 14–21.
- [6] Ali and Saleh, 2012; Borowitzka, (2013), Dependency of Microalgal Production on Biomass and the Relationship to Yield and Bioreactor Scale-up for Biofuels: a Statistical Analysis of 60+ Years of Algal Bioreactor Data Bioenerg. Res. (2017) 10:267 287DOI 10.1007/s12155-016-9787-2
- [7] Priyadarshani and Rath, (2012) Bioactive compounds from Microalgae and Cyanobacteria: Utility and Applications, IJPSR, 2012; Vol. 3(11): 4123-4130.
- [8] Pandeeey, J.P.and Tiwari, A.(2010). Optimization of Biomass Production of *Spirulina* maxima, Journal of Algal Biomass Utilization, 1(2)2032
- [9] Newsome, A. G., Culver, C. A., & Van Breemen, R. B. (2014). Nature's palette: The search for natural blue colorants. Journal of Agricultural and Food Chemistry, 62(28), 6498–6511. https://doi.org/10.1021/jf501419q.
- [10] Raja, R., Hemaiswarya, S., et al., (2007). Exploitation of Dunaliella for β-carotene production. Appl. Microbiol. Biotechnol. 74, 517–523
- [11] Tarantino LM. (2003). Agency Response Letter GRAS Notice No. GRN000127. FDA Home page, October.
- [12] Belay A. (2002). The potential application of *Spirulina (Arthrospira)* as a nutritional and therapeutic supplement in Health management. *Journal of the American Nutraceutical Association*.;5:27–48. [Google Scholar]
- [13] Khan Z, Bhadouria P, Bisen PS.(2005) Nutritional and therapeutic potential of *Spirulina*. *Current Pharmaceutical Biotechnology*.;6(5):373–379. [PubMed] [Google Scholar]
- [14] Morais MG, Stillings C, Dersch R, et al. (2010).Preparation of nanofibers containing the microalga *Spirulina* (*Arthrospira*). *Bioresource Technology*.;101(8):2872–2876. [PubMed] [Google Scholar]
- [15] Qureshi MA, Kidd NT, Ali RA. (1995) *Spirulina* platensis extract enhances chicken macrophage functions after in vitro exposure. *Journal of Nutritional Immunology*.;3 (4):35–45. [Google Scholar]
- [16] Ramier J, Bouderlique T, Stoilova O, et al. (2014). Biocomposite scaffolds based on electrospun poly(3-hydroxybutyrate) nanofibers and electrosprayed hydroxyapatite nanoparticles for bone tissue engineering applications. *Materials Science and Engineering: C*.;38:161–169. [PubMed] [Google Scholar]
- [17] Sudesh K, Abe H, Doi Y. (2000). Synthesis, structure and properties of polyhydroxyalkanoates: biological polyesters. *Progress in Polymer Science*.;25(10):1503–1555. [Google Scholar]
- [18] Jau M, Yew S, Toh PSY, et al. (2005). Biosynthesis and mobilization of poly(3-hydroxybutyrate) [P(3HB)] by *Spirulina platensis*. *International Journal of Biological Macromolecules*.;36(3):144–151. [PubMed] [Google Scholar
- [19] California Grown Earthrise Californian *Spirulina*. earthrise.com
- [20] https://www.earthrise.com > california-grown